

Polynomial hulls and positive currents

Tien-Cuong Dinh

February 1, 2008

Abstract

Let T be a positive plurisubharmonic current of bidimension (p, p) and let $\delta > 0$. Assume that the Lelong number of T satisfies $\nu(T, a) \geq \delta$ on a dense subset of $\text{supp}(T)$ (rectifiable currents satisfy this condition). Then $T = \varphi[X]$, where X is a complex subvariety of pure dimension p and φ is a weakly plurisubharmonic function on X . We have an analogous result for plurisuperharmonic currents. We also introduce and study a notion of polynomial p -hull.

1 Introduction

Let V be a complex manifold of dimension n . A current T of bidimension (p, p) or bidegree $(n - p, n - p)$ in V is called *plurisubharmonic* if $\text{dd}^c T$ is a positive current, is called *plurisuperharmonic* if $-\text{dd}^c T$ is a positive current and is called *pluriharmonic* if $\text{dd}^c T = 0$. In particular, every closed current is pluriharmonic. In this paper, we study some properties of positive plurisubharmonic and plurisuperharmonic currents.

Positive plurisubharmonic and plurisuperharmonic currents are very useful in many mathematical problems. In [17], Garnett proved that every laminated closed set supports a positive pluriharmonic currents (see also [8]). Duval and Sibony used positive plurisuperharmonic currents to describe the polynomial hull of compact sets in $\mathbb{C}^n$ [14]. Gauduchon, Harvey, Lawson, Michelson, Alessandrini, Bassanelli, etc. studied non-Kähler geometry using (smooth or not) pluriharmonic currents [1].

Many important properties of these classes of currents have been considered in the papers of Skoda [28], Sibony, Berndtsson, Fornæss [25], [8], [16],

Alessandrini and Bassanelli [2, 3, 7], etc. In particular, we know that if T is a positive plurisubharmonic current, one can define a density $\nu(T, a)$ of T at every point a . This density is called Lelong number of T at a .

We will prove that if the level set $\{\nu(T, a) \geq \delta\}$ is dense in the support of T for a suitable $\delta > 0$, then the support X of T is a complex subvariety of pure dimension p and $T = \varphi[X]$, where φ is a weakly plurisubharmonic function on X . In particular, every rectifiable positive plurisubharmonic current is closed. Moreover, it has the form $c[X]$, where X is a complex subvariety and c is essentially equal to a positive integer in each component of X . We will also prove that if T is positive pluriharmonic (resp. plurisuperharmonic) and its support has locally finite $2p$ -dimensional Hausdorff measure then the support X of T is a complex subvariety of pure dimension p and $T = \varphi[X]$, where φ is a weakly pluriharmonic (resp. plurisuperharmonic) function on X .

For closed positive currents, an analogous result has been proved by King [20] (see also [27, 26]). The proof of King does not work in the case of plurisubharmonic and plurisuperharmonic currents. Our proof relies on some recent progress on the polynomial hull of finite length compact sets [10, 11, 22]. We also use a theorem of Shiffman on separately holomorphic functions and some results of Alessandrini-Bassanelli on the Lelong number.

In order to study the currents of bidimension (p, p) we will introduce the notion of polynomial p -hull which is the usual polynomial hull when $p = 1$. We will extend the Wermer theorem for p -hull.

2 Polynomial hulls and pseudoconcave sets

Let $\Gamma \subset \mathbb{C}^n$ be a compact set. The *polynomial hull* of Γ is the compact set $\widehat{\Gamma}$ defined by the following formula

$$\widehat{\Gamma} := \{x \in \mathbb{C}^n, |P(x)| \leq \max_{z \in \Gamma} |P(z)| \text{ for any polynomial } P\}.$$

We now introduce the notions of p -pseudoconcavity and polynomial p -hull.

Let V be a complex manifold of dimension $n \geq 2$ and X a closed subset of V . Let $1 \leq p \leq n - 1$ be an integer.

Definition 1 We say that X is *p -pseudoconcave subset of V* if for every open set $U \subset\subset V$ and every holomorphic map f from a neighbourhood of $\overline{U}$ into

$\mathbb{C}^p$ we have $f(X \cap U) \subset \mathbb{C}^p \setminus \Omega$ where Ω is the unbounded component of $\mathbb{C}^p \setminus f(X \cap bU)$.

This means X has no “locally peak point” for holomorphic maps into $\mathbb{C}^p$. In particular, X satisfies the maximum principle, i.e. holomorphic functions admit no local strict maximum modul on X . Observe that if V is a submanifold of another complex manifold V' , then X is p -pseudoconcave in V if and only if X is p -pseudoconcave in V' .

By the argument principle, every complex subvariety of pure dimension p of V is p -pseudoconcave. It is clear that if X is p -pseudoconcave, the $2p$ -dimensional Hausdorff measure of X is strictly positive. If $1 \leq p \leq q \leq n-1$ and X is q -pseudoconcave then X is p -pseudoconcave. If $g : V \longrightarrow \mathbb{C}^{p-k}$ is a holomorphic map and $X \subset V$ is p -pseudoconcave then $g^{-1}(x) \cap X$ is k -pseudoconcave for every $x \in \mathbb{C}^{p-k}$. We have the following proposition.

Proposition 1 *Let T be a positive plurisuperharmonic current of bidimension (p, p) on a complex manifold V . Then the support $\text{supp}(T)$ of T is p -pseudoconcave in V .*

Proof— Since the problem is local we can assume that V is a ball of $\mathbb{C}^n$. Assume that $X := \text{supp}(T)$ is not p -pseudoconcave. Then there exist an open set $U \subset\subset V$ and a holomorphic map $f : V' \longrightarrow \mathbb{C}^p$ such that $f(X \cap U) \not\subset \mathbb{C}^p \setminus \Omega$, where V' is a neighbourhood of $\overline{U}$ and Ω is the unbounded component of $\mathbb{C}^p \setminus f(X \cap bU)$.

Let $\Phi : V' \longrightarrow \mathbb{C}^p \times V'$ the holomorphic map given by $\Phi(z) := (f(z), z)$. Choose a bounded domain U' in $\mathbb{C}^p \times V'$ such that $U' \cap \Phi(V') = \Phi(U)$. Set $X' := \Phi(X \cap V')$ and $T' := 1_{U'} \Phi_*(T)$. Then T' is a positive plurisuperharmonic current in U' . Let $\pi : \mathbb{C}^{n+p} \longrightarrow \mathbb{C}^p$ be the linear projection on the first p coordinates. We have $\pi(\overline{X'} \cap bU') = f(X \cap bU)$ and $\pi(X' \cap U') = f(X \cap U) \not\subset \mathbb{C}^p \setminus \Omega$. The open set Ω is also the unbounded component of $\mathbb{C}^p \setminus \pi(\overline{X'} \cap bU')$.

Therefore $\pi_*(T')$ defines a positive plurisuperharmonic current in Ω which is null in $\mathbb{C}^p \setminus \pi(\overline{X'})$. Hence there is a positive plurisuperharmonic function ψ on Ω , null on $\mathbb{C}^p \setminus \pi(\overline{X'})$ such that $\pi_*(T') = \psi[\Omega]$ in Ω . By plurisuperharmonicity, this function is identically null. Fix a small open $W \subset \mathbb{C}^{n+p}$ such that $W \cap X' \neq \emptyset$ and $\pi(\overline{W}) \subset \Omega$. Set $\Psi := (idz_1 \wedge d\overline{z}_1 + \cdots + idz_p \wedge d\overline{z}_p)^p$. Then the positive measure $T' \wedge \pi^*(\Psi)$ is null in W since its push-forward by π is null. This still holds for every small linear pertubation π_ϵ of π . On

the other hand, we can construct a strictly positive (p, p) -form ψ of $\mathbb{C}^{n+p}$ as a linear combinaison of $\pi_e^*(\Psi)$. Then we have $\langle T', 1_W \psi \rangle = 0$. Since T' is positive, $T' = 0$ on W . This is a contradiction.

□

Remark 1 We can also prove Proposition 1 by using the Oka inequality of Fornæss and Sibony [16]

Proposition 2 *Let Γ be a compact subset of $\mathbb{C}^n$. Denote by $\mathcal{F}$ the family of p -pseudoconcave subsets of $\mathbb{C}^n \setminus \Gamma$ which are bounded in $\mathbb{C}^n$. Then the union Σ of elements of $\mathcal{F}$ is also belong to $\mathcal{F}$ (then Σ is the biggest element of $\mathcal{F}$).*

Proof— Let $S_1, S_2, \dots$ be elements of $\mathcal{F}$ such that $S_i \subset S_{i+1}$. We show that $S := \overline{\bigcup S_i}$ belongs to $\mathcal{F}$. By maximum principle, $S_i \subset \widehat{\Gamma}$. Then S is bounded in $\mathbb{C}^n$.

Assume that S does not belong to $\mathcal{F}$. Then there are a point $p \in S$, a neighbourhood $U \subset \subset \mathbb{C}^n \setminus \Gamma$ of p and a holomorphic map f from a neighbourhood of $\overline{U}$ into $\mathbb{C}^p$ such that $f(p)$ belongs to the unbounded component Ω of $\mathbb{C}^p \setminus f(S \cap bU)$. Fix small neighbourhoods V of p and W of $S \cap bU$ such that $f(\overline{V})$ is included in the unbounded component of $\mathbb{C}^p \setminus f(\overline{W})$.

For i big enough, we have $S_i \cap bU \subset W$ and $S_i \cap V \neq \emptyset$. Moreover, if p is a point in $S_i \cap V$, $f(p)$ belongs to the unbounded component of $\mathbb{C}^p \setminus f(S_i \cap bU)$. This contradicts the p -pseudoconcavity. Thus $S \in \mathcal{F}$.

Now, by Zorn axiom, we can choose an element Σ' of $\mathcal{F}$ which is maximal for the inclusion. If $\Sigma' \neq \Sigma$ then there is an element $S \in \mathcal{F}$ such that $S \not\subset \Sigma'$. The set $S \cup \Sigma'$ is p -pseudoconcave in $\mathbb{C}^n \setminus \Gamma$ and it belongs to $\mathcal{F}$. This contradicts the maximality of Σ' .

□

Definition 2 Let Γ and Σ be as in Proposition 2. We say that $\Sigma \cup \Gamma$ is the *polynomial p -hull* of Γ and we denote it by $\text{hull}(\Gamma, p)$.

The following proposition shows that we obtain the usual polynomial hull when $p = 1$.

Proposition 3 *We have $\text{hull}(\Gamma, 1) = \widehat{\Gamma}$ for every compact subset Γ of $\mathbb{C}^n$.*

Proof— By maximum principle, we have $\text{hull}(\Gamma, 1) \subset \widehat{\Gamma}$. Now let $z \in \widehat{\Gamma} \setminus \Gamma$, we will prove that $z \in \text{hull}(\Gamma, 1)$. By Duval-Sibony theorem [14], there are a

positive current T of bidimension $(1, 1)$ with compact support in $\mathbb{C}^n$ and a measure μ with support in Γ such that $\text{dd}^c T = \mu - \delta_z$ where δ_z is the Dirac mass at z . By Proposition 1, $\text{supp}(T)$ is 1-pseudoconcave in $\mathbb{C}^n \setminus \Gamma$. Thus $\text{supp}(T) \subset \text{hull}(\Gamma, 1)$ and $z \in \text{hull}(\Gamma, 1)$. □

3 Hulls of finite Hausdorff measure sets

In this section, we first recall some results on the polynomial hull of finite length compact sets and then we will extend them for polynomial p -hull.

Denote by $\mathcal{H}^k$ the Hausdorff measure of dimension k . A compact set $\Gamma \subset \mathbb{C}^n$ is called *geometrically k -rectifiable* if $\mathcal{H}^k(\Gamma)$ is finite and the geometric tangent cone of Γ is a real space of dimension k at $\mathcal{H}^k$ -almost every point in Γ . If a compact set Γ is geometrically k -rectifiable, it is $(\mathcal{H}^k, k)$ -*rectifiable* [23, p.208], i.e. there exist $\mathcal{C}^1$ manifolds $V_1, V_2, \dots$ such that $\mathcal{H}^k(\Gamma \setminus \cup V_m) = 0$ [15, 3.1.16, 3.2.18]. Now, suppose that $V_1, V_2, \dots$ are $\mathcal{C}^1$ oriented manifolds of dimension k in $\mathbb{C}^n$, $K_i \subset V_i$ are compact sets and $n_1, n_2, \dots$ are integers such that $\sum |n_i| \mathcal{H}^k(K_i) < +\infty$. Then we can define a current S of dimension k by

$$\langle S, \psi \rangle := \sum n_i \int_{K_i} \psi$$

for any test form ψ of degree k having compact support. Such a current is called *rectifiable current*.

We have the following theorem.

Theorem 1 *Let Γ be a compact subset of $\mathbb{C}^n$. Assume that Γ is geometrically $(2p-1)$ -rectifiable with $1 \leq p \leq n-1$. Then $\text{hull}(\Gamma, p) \setminus \Gamma$ is a complex subvariety of pure dimension p (possibly empty) of $\mathbb{C}^n \setminus \Gamma$. Moreover, $\text{hull}(\Gamma, p) \setminus \Gamma$ has finite $2p$ -dimensional Hausdorff measure and the boundary of the integration current $[\text{hull}(\Gamma, p) \setminus \Gamma]$ is a rectifiable current of dimension $2p-1$ and has multiplicity 0 or 1 $\mathcal{H}^{2p-1}$ -almost everywhere on Γ .*

The last property of the current $[\text{hull}(\Gamma, p) \setminus \Gamma]$ is called *Stokes formula*. In the case $p = 1$, this theorem is proved in [10, 11, 22] and it generalizes the results of Wermer [30], Bishop [9], Stolzenberg [29], Alexander [4] and Harvey-Lawson [18] (see also [13]).

In order to prove Theorem 1, we will slice Γ by complex planes of dimension $n - p + 1$ and we will apply the known result in the case $p = 1$. We will also use a theorem of Shiffman on separately holomorphic functions.

Set $X := \text{hull}(\Gamma, p) \setminus \Gamma$. Observe that if L is a linear complex $(n - p + 1)$ -plane then $X \cap L$ is 1-pseudoconcave in $L \setminus \Gamma$. Thus $X \cap L$ is included in the polynomial hull of $\Gamma \cap L$. Moreover, by Sard theorem, for almost every L the intersection $\Gamma \cap L$ is geometrically 1-rectifiable (see also [10]). Therefore $\text{hull}(\Gamma \cap L, 1) \setminus \Gamma$ is a complex subvariety of pure dimension 1 of $L \setminus \Gamma$. We need the following lemma.

Lemma 1 *Let V be a complex manifold of dimension $n \geq 2$ and $X \subset V$ a p -pseudoconcave subset. If X is included in a complex subvariety Σ of pure dimension p of V , then X is itself a complex subvariety of pure dimension p of V .*

Proof— Recall that every non empty p -pseudoconcave set has positive $\mathcal{H}^{2p}$ measure. Assume that X is not a complex subvariety of pure dimension p of V . Then there is a point $a \in X$ such that a is a regular point of Σ and $a \in \overline{\Sigma \setminus X}$. Let $V' \subset V$ be a small ball of centre a satisfying $V' \cap \Sigma \setminus X \neq \emptyset$. Since V' is small, we can choose a projection $\pi : V' \rightarrow \mathbb{C}^p$ such that π is injective on $\Sigma \cap V'$. Then $\pi(a)$ is belong to the unbounded component of $\mathbb{C}^p \setminus \pi(X \cap bW)$ for every neighbourhood $W \subset\subset V'$ of a . This is impossible. $\square$

By the latter lemma, for almost every L (such that $\Gamma \cap L$ is geometrically 1-rectifiable), $X \cap L$ is a complex subvariety of pure dimension 1 of $L \setminus \Gamma$. Let E be a complex $(n - p)$ -plane which does not meet Γ . By Sard theorem, for almost every L passing through E , the intersection $\Gamma \cap L$ is geometrically 1-rectifiable. We deduce that $X \cap E = (X \cap L) \cap E$ is a finite set. We now use a Shiffman theorem on separately holomorphic functions in order to complete the proof of Theorem 2.

Let a be a point of X . We show that X is a complex variety of pure dimension p in a neighbourhood of a . Indeed, choose a coordinates system such that $\Pi(a) \notin \Pi(\Gamma)$, where $\Pi : \mathbb{C}^n \rightarrow \mathbb{C}^p$ is the linear projection given by $\Pi(z) := (z_1, \dots, z_p)$. Set $\pi_i : \mathbb{C}^n \rightarrow \mathbb{C}^{p-1}$, $\pi_i(z) := (z_1, \dots, z_{i-1}, z_{i+1}, \dots, z_p)$. Using a linear change of coordinates, we may suppose without loss of generality that for $\mathcal{H}^{2p-2}$ -almost every $x \in \mathbb{C}^{p-1}$ and for every $1 \leq i \leq n$, $X \cap \pi_i^{-1}(x)$ is a complex subvariety of pure dimension 1 of $\pi_i^{-1}(x) \setminus \Gamma$.

Fix a small open neighbourhood V of $\Pi(a)$ such that $\Pi^{-1}(V) \cap \Gamma = \emptyset$. Put $z = (z', z'')$, $z' := (z_1, \dots, z_p)$ and $z'' := (z_{p+1}, \dots, z_n)$. Let g be a function which is defined in a subset V' of total measure of V . This function is called *separately holomorphic* if for every $1 \leq i \leq p$ and $\mathcal{H}^{2p-2}$ -almost every $x \in \mathbb{C}^{p-1}$ the restriction of g on $V' \cap \pi_i^{-1}(x)$ can be extended to a holomorphic function on $V \cap \pi_i^{-1}(x)$. For any $k \geq 0$ and $p+1 \leq m \leq n$, we define the following measurable function on V

$$g_{m,k}(z') := \sum_{z \in \Pi^{-1}(z') \cap X} z_m^k.$$

The function

$$g_0(z') := g_{m,0}(z) = \#(\Pi^{-1}(z') \cap X)$$

is independent on m and takes only positive integer values.

By the choice of coordinates, $g_{m,k}(z')$ is $\mathcal{H}^{2p}$ -almost everywhere equal to a separately holomorphic function. Thanks to Shiffman Theorem [24], $g_{m,k}(z')$ is equal $\mathcal{H}^{2p}$ -almost everywhere to a function $\tilde{g}_{m,k}(z')$ which is holomorphic on V . In particular, $\tilde{g}_{m,0}(z')$ is equal to an integer r which does not depend on m and z' .

Now, consider the following equation system:

$$\sum_{i=1}^r [z_m^{(i)}]^k = \tilde{g}_{m,k}(z') \quad \text{with } p+1 \leq m \leq n \text{ and } 1 \leq k \leq r.$$

The set of points $(z', z_{p+1}^{(i_{p+1})}, \dots, z_n^{(i_n)})$ given by solutions of the system above is a complex subvariety Σ of pure dimension p of $\Pi^{-1}(V)$. We have $X \cap \Pi^{-1}(x) \subset \Sigma$ for $\mathcal{H}^{2p}$ -almost every $x \in V$. We show that $X \cap \Pi^{-1}(V) \subset \Sigma$. Assume this is not true. Choose a point $b \in X \cap \Pi^{-1}(V) \setminus \Sigma$ and a neighbourhood $W \subset \subset \Pi^{-1}(V) \setminus \Sigma$ of b . The set $\Pi(X \cap \overline{W})$ has zero $\mathcal{H}^{2p}$ measure. But for almost every complex $(n-p+1)$ -plane L passing through $\Pi^{-1}(\Pi(b))$, the intersection $X \cap L$ is a complex subvariety of pure dimension 1 of $L \setminus \Gamma$ which contains b . This implies that the set $\Pi(X \cap \overline{W})$ has positive $\mathcal{H}^{2p}$ measure. We reach a contradiction.

Now we have already know that $X \cap \Pi^{-1}(V) \subset \Sigma$. By Lemma 1, $X \cap \Pi^{-1}(V)$ is a complex subvariety of pure dimension p of $\Pi^{-1}(V)$. Then X is a complex subvariety of pure dimension p of $\mathbb{C}^n \setminus \Gamma$. The rest of the proof follows along the same lines as the proof of Stokes formula given in [21, 10, 11].

□

When Γ is included in the boundary of a smooth convex domain, its rectifiability in Theorem 1 is non necessary. We have the following result.

Theorem 2 *Let D be a strictly convex domain with $\mathcal{C}^2$ boundary of $\mathbb{C}^n$. Let Γ be a compact set in the boundary of D . Assume that the $(2p - 1)$ -dimensional Hausdorff measure of Γ is finite. Then $\text{hull}(\Gamma, p) \setminus \Gamma$ is a complex subvariety of pure dimension p (possibly empty) of $\mathbb{C}^n \setminus \Gamma$. Moreover, $\text{hull}(\Gamma, p) \setminus \Gamma$ has finite $2p$ -dimensional Hausdorff measure and the boundary of the integration current $[\text{hull}(\Gamma, p) \setminus \Gamma]$ is a rectifiable current of dimension $2p - 1$ and has multiplicity 0 or 1 $\mathcal{H}^{2p-1}$ -almost everywhere on Γ .*

In the case $p = 1$, this result is proved by Lawrence [22] as a corollary of Theorem 1. In order to apply Theorem 1, Lawrence show that if a compact set $\Gamma' \subset \Gamma$ is minimal with respect to the property that its polynomial hull contains some point $w \in D$, then it is geometrically 1-rectifiable. Using this result of Lawrence, the proof of Theorem 1 remains valid for Theorem 2.

Remark 2 Under the hypothesis of Theorems 1 and 2, by Lemma 1, every p -pseudoconcave subset of $\mathbb{C}^n \setminus \Gamma$ which is bounded in $\mathbb{C}^n$, is a complex subvariety of pure dimension p of $\mathbb{C}^n \setminus \Gamma$ and is included in X . By Proposition 1, every positive current T with compact support satisfying $\text{dd}^c T \leq 0$ in $\mathbb{C}^n \setminus \Gamma$ has the form $\varphi[X]$, where φ is a weakly plurisuperharmonic function in X (see the next section).

Corollary 1 *Let V be a complex manifold of dimension $n \geq 2$ and X a p -pseudoconcave subset of V . Let K be a compact subset of V which admits a Stein neighbourhood. Assume that the $2p$ -dimensional Hausdorff measure of $X \setminus K$ is locally finite in $V \setminus K$. Then X is a complex subvariety of pure dimension p of V .*

Proof— Fix a point $a \in X \setminus K$ and $B \subset\subset V \setminus K$ an open ball contening a such that $X \cap bB$ has finite $\mathcal{H}^{2p-1}$ measure. Then $X \cap B$ is p -pseudoconcave in B . We can consider B as an open ball of $\mathbb{C}^n$. Then $X \cap B \subset \text{hull}(X \cap bB, p)$. By Remark 2, $X \cap B$ is a complex subvariety of pure dimension p of B . This implies that $X \setminus K$ is a complex subvariety of pure dimension p of $V \setminus K$.

Since we can replace V by a Stein neighbourhood of K , suppose without loss of generality that V is a submanifold of $\mathbb{C}^N$. Observe that X is also p -pseudoconcave in $\mathbb{C}^N$. Let B' be a ball containing the compact K such that

$X \cap B'$ has finite $\mathcal{H}^{2p-1}$ measure. Then we obtain $X \cap B' \subset \text{hull}(X \cap bB', p)$. By Remark 2, $X \cap B'$ is a complex subvariety of pure dimension p of B' . This completes the proof. $\square$

4 Positive currents

Let V be a complex manifold of dimension n . Let X be a complex subvariety of pure dimension p of V . A function $\varphi \in L^1_{\text{loc}}(X)$ is called *weakly plurisubharmonic* if it is essentially bounded from above and $\text{dd}^c(\varphi[X])$ is a positive current. In the regular part of X , φ is equal almost everywhere to an usual plurisubharmonic function. We say that φ is *weakly plurisuperharmonic* if $-\varphi$ is weakly plurisubharmonic and that φ is *weakly pluriharmonic* if both φ and $-\varphi$ are weakly plurisubharmonic.

If φ is a positive weakly plurisubharmonic function on X , we can define a positive plurisubharmonic current $T := \varphi[X]$ as follows:

$$\langle T, \psi \rangle := \int_X \varphi \psi$$

for every test form ψ of bidegree (p, p) compactly supported in V . According to Bassanelli [7, 1.24, 4.10], every positive plurisubharmonic current with support in X can be obtained by this way. This is also true for positive plurisuperharmonic and pluriharmonic currents.

Now, let T be a positive plurisubharmonic current of bidimension (p, p) of V . Thanks to a Jensen type formula [3, 28, 12], if V is an open set of $\mathbb{C}^n$ which contains a ball $B(a, r_0)$, the function

$$\frac{1}{2^p \pi^p r^{2p}} \langle T, 1_{B(a, r)} \omega^p \rangle, \text{ for } 0 \leq r < r_0,$$

is non-decreasing, where $\omega := idz_1 \wedge d\bar{z}_1 + \cdots + idz_n \wedge d\bar{z}_n$. Therefore, we can define

$$\nu(T, a) := \lim_{r \rightarrow 0} \frac{1}{2^p \pi^p r^{2p}} \langle T, 1_{B(a, r)} \omega^p \rangle.$$

This limit is called *Lelong number* of T at the point a . In [3], Alessandrini and Bassanelli proved that $\nu(T, a)$ does not depend on coordinates. Thus, the Lelong number is well defined for every manifold V .

We have the following theorem.

Theorem 3 *Let T be a positive plurisubharmonic current of bidimension (p, p) in a complex manifold V of dimension n . Assume that there exists a real number $\delta > 0$ such that the level set $\{z \in V, \nu(T, z) \geq \delta\}$ is dense in the support $\text{supp}(T)$ of T . Then $\text{supp}(T)$ is a complex subvariety of pure dimension p of V and there exists a weakly plurisubharmonic function φ on $X := \text{supp}(T)$ such that $T = \varphi[X]$.*

Remark 3 In the case of closed positive currents, an analogous result is proved by King [20].

It is sufficient to prove that the support $X = \text{supp}(T)$ of T is a complex subvariety of pure dimension p of V . Since the problem is local, we can suppose that V is a ball of $\mathbb{C}^n$. By Corollary 1, we have to prove that X has locally finite $\mathcal{H}^{2p}$ measure and X is p -pseudoconcave. Consider now the trace measure $\sigma := T \wedge \omega^p$ of T .

Proposition 4 *Let T be a plurisubharmonic current of bidimension (p, p) in a complex manifold V . Then for every $\delta > 0$ the level set $\{\nu(T, a) \geq \delta\}$ is closed and has locally finite $\mathcal{H}^{2p}$ measure. Moreover, under the hypothesis of Theorem 3, we have $\nu(T, a) \geq \delta$ for every $a \in X$ and X has locally finite $\mathcal{H}^{2p}$ measure.*

Proof— We may suppose without loss of generality that V is an open subset of $\mathbb{C}^n$. Set $Y := \{\nu(T, a) \geq \delta\}$. Let $(a_n) \subset Y$ be a sequence which converges to a point $a \in V$. Fix an $r > 0$ such that $B(a, r) \subset V$. We have

$$\sigma(B(a, r)) \geq \sigma(B(a_n, r - |a_n - a|)) \geq 2^p \pi^p \delta (r - |a_n - a|)^{2p},$$

which implies that

$$\sigma(B(a, r)) \geq 2^p \pi^p \delta r^{2p}$$

and

$$\nu(T, a) = \lim_{r \rightarrow 0} \frac{\sigma(B(a, r))}{2^p \pi^p r^{2p}} \geq \delta.$$

Therefore, $a \in Y$ and hence Y is closed. The last inequality also implies that Y has locally finite $\mathcal{H}^{2p}$ measure [15, 2.10.19(3)].

□

Lemma 2 *Let T be a positive current of bidimension (p, p) in $\mathbb{C}^n$ with compact support. Let $\pi : \mathbb{C}^n \longrightarrow \mathbb{C}^p$ be the linear projection on the first p coordinates, $a \in \mathbb{C}^n$ and $b := \pi(a)$. Assume that T is plurisubharmonic on $\pi^{-1}(B(b, r))$ for some $r > 0$. Then $\langle T, 1_{\pi^{-1}(B(b, r))} \Psi \rangle \geq 2^p \pi^p r^{2p} \nu(T, a)$, where $\Psi := (\text{id} z_1 \wedge d\bar{z}_1 + \cdots + \text{id} z_p \wedge d\bar{z}_p)^p$. In particular, the Lelong number of $\pi_*(T)$ at b is greater or equal to $\nu(T, a)$.*

Proof— We may suppose without loss of generality $a = 0$ and $b = 0$. Set $z' := (z_1, \dots, z_p)$, $z'' := (z_{p+1}, \dots, z_n)$, $\omega := \text{id} z_1 \wedge d\bar{z}_1 + \cdots + \text{id} z_p \wedge d\bar{z}_p$ and fix an $\epsilon > 0$. Let $\Phi : \mathbb{C}^n \longrightarrow \mathbb{C}^n$ be the linear map given by $\Phi_\epsilon(z) := (z', \epsilon z'')$ and set $T' := (\Phi_\epsilon)_* T$. Since the Lelong number is independant on coordinates [3], we have $\nu(T', 0) = \nu(T, 0)$. By Jensen type formula we have

$$\langle T', 1_{\pi^{-1}(B(0, r))} \omega^p \rangle \geq 2^p \pi^p r^{2p} \nu(T', 0)$$

since $\pi^{-1}(B(0, r))$ contains the ball $B(0, r)$ of $\mathbb{C}^n$. From this inequality, it follows that

$$\langle T, 1_{\pi^{-1}(B(0, r))} \Phi_\epsilon^*(\omega^p) \rangle \geq 2^p \pi^p r^{2p} \nu(T, 0).$$

On the other hand, when $\epsilon \rightarrow 0$, $\Phi_\epsilon^*(\omega^p)$ converges uniformly to Ψ . Thus,

$$\langle T, 1_{\pi^{-1}(B(0, r))} \Psi \rangle \geq 2^p \pi^p r^{2p} \nu(T, 0).$$

This implies

$$\nu(\pi_*(T), 0) = \lim_{r \rightarrow 0} \frac{1}{2^p \pi^p r^{2p}} \langle T, 1_{\pi^{-1}(B(0, r))} \Psi \rangle \geq \nu(T, 0),$$

which completes the proof. □

The following lemma completes the proof of Theorem 3.

Lemma 3 *The set X is p -pseudoconcave.*

Proof— Assume that X is not p -pseudoconcave. Then there exist an open set $U \subset\subset V$ and a holomorphic map $f : V' \longrightarrow \mathbb{C}^p$ such that $f(X \cap U) \not\subset \mathbb{C}^p \setminus \Omega$, where V' is a neighbourhood of $\bar{U}$ and Ω is the unbounded component of $\mathbb{C}^p \setminus f(X \cap bU)$.

Consider $\Phi : V' \longrightarrow \mathbb{C}^p \times V'$ the holomorphic map given by $\Phi(z) := (f(z), z)$. We next choose a domain $U' \subset\subset \mathbb{C}^p \times V'$ such that $\Phi(U)$ is a

submanifold of U' . Then $T' := 1_{U'}\Phi_*(T)$ is a positive plurisubharmonic current in U' . Moreover it is easy to see that $\nu(T', a) \geq \delta'$ for some $\delta' > 0$ and for every $a \in X' := \text{supp}(T')$. Let $\pi : \mathbb{C}^{p+n} \rightarrow \mathbb{C}^p$ be the linear projection on the first p coordinates. We have $\pi(\overline{X}' \cap bU') = f(X \cap bU)$ and $\pi(X' \cap U') = f(X \cap U) \not\subset \mathbb{C}^p \setminus \Omega$. The open set Ω is also the unbounded component of $\mathbb{C}^p \setminus \pi(\overline{X}' \cap bU')$.

Therefore $\pi_*(T')$ defines a positive plurisubharmonic current in Ω which is null in $\mathbb{C}^p \setminus \pi(\overline{X}')$. Hence there is a positive plurisubharmonic function ψ on Ω , null on $\mathbb{C}^p \setminus \pi(\overline{X}')$ such that $\pi_*(T') = \psi[\Omega]$ in Ω . Thus for every $x \in \Omega$ we have $\nu(\pi_*(T'), x) = \psi(x)$. By Lemma 2, we have $\psi(x) \geq \delta'$ for every $x \in \pi(X') \cap \Omega$.

Now fix a point x in the boundary Σ of $\pi(X')$ in Ω . Let $c \in \Omega$ be a point close to x . Let $b \in \pi(X')$ such that $\text{dist}(b, c) = \text{dist}(\pi(X'), c)$. Since c is close to x , we have $b \in \Sigma$ and $B(c, |b - c|) \subset \Omega$. By submean property of plurisubharmonic functions and the fact that $\Psi = 0$ on $B(c, |b - c|)$, we obtain

$$\psi(b) = \lim_{r \rightarrow 0} \frac{1}{2^p \pi^p r^{2p}} \int_{B(b, r)} \psi(z) \Psi = \lim_{r \rightarrow 0} \frac{1}{2^p \pi^p r^{2p}} \int_{B(b, r) \setminus B(c, |b - c|)} \psi(z) \Psi.$$

By uppersemicontinuity, the last limit is smaller or equal to

$$\lim_{r \rightarrow 0} \frac{1}{2^p \pi^p r^{2p}} \int_{B(b, r) \setminus B(c, |b - c|)} \psi(b) \Psi = \frac{\psi(b)}{2}.$$

Thus $\psi(b) \leq \frac{1}{2}\psi(b)$ and $\psi(b) \leq 0$. This contradicts the fact that $\varphi(b) \geq \delta'$. $\square$

Corollary 2 *Let T be a positive plurisubharmonic current of bidimension (p, p) in a complex manifold V of dimension n . If T is locally rectifiable, then there exists a locally finite family of complex subvarieties $(X_i)_{i \in I}$ of pure dimension p of V and positive integers n_i such that $T = \sum_{i \in I} n_i [X_i]$. In particular, T is closed.*

Remark 4 King has proved the same result for rectifiable closed positive current [20]. Harvey, Shiffman and Alexander [5, 19] proved it for rectifiable closed currents (in this case, the positivity is not necessary).

Proof— By [15, 4.1.28(5)], $\nu(T, a)$ is a strictly positive integer for a in a dense subset of $X := \text{supp}(T)$. This, combined with Theorem 3, implies that X is a complex subvariety of pure dimension p of V and $T = \varphi[X]$, where φ is a weakly plurisubharmonic function on X . Since T is rectifiable, the function φ has integer values $\mathcal{H}^{2p}$ -almost everywhere. Therefore φ is essentially equal to a positive integer in each irreducible component of X . $\square$

The following theorem is a variant of Theorem 3 (see also Example 1).

Theorem 3' *Let T be a positive plurisubharmonic current of bidimension (p, p) in a complex manifold V of dimension n . Assume that $\text{supp}(T)$ has locally finite $\mathcal{H}^{2p}$ measure and the set $\mathcal{E} := \{z \in \text{supp}(T), \nu(T, z) = 0\}$ has zero $\mathcal{H}^{2p-1}$ measure. Then $\text{supp}(T)$ is a complex subvariety of pure dimension p of V and there exists a weakly plurisubharmonic function φ on $X := \text{supp}(T)$ such that $T = \varphi[X]$.*

Proof— We only need to prove Lemma 3. More precisely, we have to find a point $b \in \Sigma \setminus \pi(\mathcal{E})$ such that the upper-density

$$\overline{\Theta}(b) := \limsup_{r \rightarrow 0} \frac{1}{2^p \pi^p r^{2p}} \int_{B(b, r) \cap \Omega} \Psi$$

of Ω at b is strictly positive. Following [31, 5.8.5, 5.9.5], the set $\Sigma' := \{b \in \Sigma, \overline{\Theta}(b) > 0\}$ has positive $\mathcal{H}^{2p-1}$ measure. It is sufficient to take b in $\Sigma' \setminus \pi(\mathcal{E})$. The last set is not empty since $\mathcal{H}^{2p-1}(\mathcal{E}) = 0$. $\square$

For plurisuperharmonic currents, we have the following theorem.

Theorem 4 *Let T be a positive plurisuperharmonic current of bidimension (p, p) in a complex manifold V of dimension n . Let K be a compact subset of V which admits a Stein neighbourhood. Assume that in $V \setminus K$ the support $\text{supp}(T)$ of T has locally finite $2p$ -dimensional Hausdorff measure. Then $\text{supp}(T)$ is a complex subvariety of pure dimension p of V and there exists a weakly plurisuperharmonic function φ on $X := \text{supp}(T)$ such that $T = \varphi[X]$. Moreover, if T is pluriharmonic, then φ is weakly pluriharmonic on X .*

Proof— By Proposition 1, $X := \text{supp}(T)$ is p -pseudoconcave subset of V . Moreover, by Corollary 1, X is a complex subvariety of pure dimension p of V . The theorem follows. $\square$

We remark here that Theorem 4 is false for positive plurisubharmonic currents. Consider an example.

Example 1 Let ψ be a positive subharmonic function in $\mathbb{C}$ which is null in the unit disk $B(0, 1)$. We can take for example $\psi(z) := \log |z|$ if $|z| > 1$ and $\psi(z) := 0$ if $|z| \leq 1$. Let f be a holomorphic function in $\mathbb{C} \setminus B(0, 1/2)$ which cannot be extended to a meromorphic function on $\mathbb{C}$. Denote by $Y \subset \mathbb{C}^2$ the graph of f over $\mathbb{C} \setminus B(0, 1/2)$. Let φ be the subharmonic function on Y given by $\varphi(z) := \psi(z_1)$. Then $T := \varphi[Y]$ is a positive plurisubharmonic current of $\mathbb{C}^2$. Its support $X \subset Y$ is not a complex subvariety of $\mathbb{C}^2$ and cannot be extended to a complex subvariety. We can construct more complicate examples by taking countables combinations of such currents.

Corollary 3 *Under the hypothesis of Theorem 3, Theorem 3' or Theorem 4, if V is compact, then there are complex subvarieties $X_1, \dots, X_k$ of pure dimension p of V and positive real numbers $c_1, \dots, c_k$ such that $T = c_1[X_1] + \dots + c_k[X_k]$. In particular, T is closed.*

Proof— By Theorems 3, 3' and 4, X is a complex subvariety of pure dimension p . Since V is compact, X has a finite number of irreducible components and φ is essentially constant in each component. □

5 Further remarks

Denote by Δ the closed unit disk of the complex line. We say that V satisfies the *p-Kontinuitätssatz* if for any smooth family of p -dimensional closed holomorphic polydiscs Δ_t^p in V indexed by $t \in [0, 1)$ such that $\cup b\Delta_t^p$ lies on a compact subset of V , then $\cup \Delta_t^p$ lies on a compact subset of V . The following definition of p -pseudoconcavity does not change the main results of this article.

Definition 1' A closed subset X of a complex manifold V is called *p-pseudoconcave* if $V \setminus X$ satisfies the local *p-Kontinuitätssatz*, i.e. for every $a \in X$ there is a neighbourhood U of a such that $U \setminus X$ satisfies the *p-Kontinuitätssatz*.

Let Γ be an oriented compact real submanifold of dimension $2p - 1$ of $\mathbb{C}^n$.

Alexander and Wermer [6] introduced an integer number $\text{link}(\Gamma, H)$ (called linking number) which depends on Γ and on a complex subvariety H of dimension $n - p$ such that $H \cap \Gamma = \emptyset$. One can define the hull of Γ by

$$\text{hull}^*(\Gamma, p) := \Gamma \cup \{z \in \mathbb{C}^n \setminus \Gamma, \text{link}(\Gamma, H) \neq 0 \text{ for every } H \text{ containing } z\}.$$

Theorem 1 is valid for this hull but the definition cannot be extended for every compact set Γ . We now try to give an analogous definition.

Let $\Gamma \subset \mathbb{C}$ be a compact set and $\Pi : \mathbb{C}^n \longrightarrow \mathbb{C}^p$ be a polynomial map. Denote by Ω the unbounded component of $\mathbb{C}^p \setminus \Pi(\Gamma)$ and set $\Gamma_\Pi := \Pi^{-1}(\mathbb{C}^p \setminus \Omega)$. Then we can define

$$\text{hull}^{**}(\Gamma, p) := \bigcap_{\Pi} \Gamma_\Pi.$$

Observe that $\text{link}(\Gamma, \Pi^{-1}(\xi)) = 0$ if Γ is an oriented compact real manifold of dimension $2p - 1$ and $\xi \in \Omega$. Moreover we have $\text{hull}(\Gamma, p) \subset \text{hull}^{**}(\Gamma, p)$ and $\widehat{\Gamma} = \text{hull}(\Gamma, 1) = \text{hull}^{**}(\Gamma, 1)$. If $\Gamma \cap L$ is a closed irreducible curve with finite length for almost every complex $(n - p + 1)$ -plane L , then by slicing method we can prove that $\text{hull}(\Gamma, p) = \text{hull}^{**}(\Gamma, p)$. In general, we do not know when $\text{hull}(\Gamma, p) = \text{hull}^{**}(\Gamma, p)$. There are different problems which are related to this question, for example: find topology properties of $\text{hull}(\Gamma, p)$, study $\text{hull}(\Gamma, p)$ when Γ is a compact CR-manifolds, describe $\text{hull}(\Gamma, p)$ using positive currents... We also hope that some notions introduced in this article can be extended and investigated in the case of almost complex manifolds.

About positive plurisubharmonic currents, the description of the set $\{\nu(T, a) > 0\}$ is far to be complete. Assume that $\nu(T, a) \geq \delta > 0$ for every a such that $\nu(T, a) \neq 0$; we do not know if A is an analytic set.

Acknowledgments. I would like to thank the Alexander von Humboldt foundation for its support, Professor Jürgen Leiterer and Dr. George Marinescu for their great hospitality and Dr. Nguyen Viet Anh for many help during the preparation of this paper.

References

- [1] *L. Alessandrini and G. Bassanelli*, Positive $\partial\bar{\partial}$ -closed currents and non-Kähler geometry, *J. Geometric Analysis*, **2** (1992), 291-316.

- [2] *L. Alessandrini and G. Bassanelli*, Plurisubharmonic currents and their extension across analytic subsets, *Forum Math.*, **5** (1993), 577-602.
- [3] *L. Alessandrini and G. Bassanelli*, Lelong numbers of positive plurisubharmonic currents, *Results in Mathematics*, **30** (1996), 191-224.
- [4] *H. Alexander*, Polynomial approximation and hulls in sets of finite linear measure in $\mathbb{C}^n$, *Amer. J. Math.* **93** (1971), 65-74.
- [5] *H. Alexander*, Holomorphic chains and the support hypothesis conjecture, *J. Am. Math. Soc.*, **10**, No.1 (1997), 123-138.
- [6] *H. Alexander and J. Wermer*, Linking numbers and boundaries of varieties, *Ann. Math. (2)*, **151**, No.1 (2000), 125-150.
- [7] *G. Bassanelli*, A cutt-off theorem for plurisubharmonic currents, *Forum Math.*, **6** (1994), 567-595.
- [8] *B. Berndtsson and N. Sibony*, The $\bar{\partial}$ equation on a positive current, *Inv. math.*, **147** (2002), 371-428.
- [9] *E. Bishop*, Analyticity in certain Banach algebras, *Trans. Amer. Math. Soc.*, **102**(1962), 507-544.
- [10] *T.C. Dinh*, Enveloppe polynomiale d'un compact de longueur finie et chaînes holomorphes à bord rectifiable, *Acta Math.*, **180**, No.1 (1998), 31-67.
- [11] *T.C. Dinh*, Orthogonal measures on the boundary of a Riemann surface and polynomial hull of compacts of finite length, *J. Funct. Anal.*, **157**, No.2 (1998), 624-649.
- [12] *J.P. Demailly*, Formules de Jensen en plusieurs variables et application arithmétique, *Bull. Soc. Math. France*, **110** (1982), 75-102.
- [13] *P. Dolbeault and G. Henkin*, Chaînes holomorphes de bord donné dans $\mathbb{CP}^n$, *Bull. Soc. Math. de France*, **125** (1997), 383-445.
- [14] *J. Duval and N. Sibony*, Polynomial convexity, rational convexity, and currents, *Duke Math. J.*, **79**, No.2 (1995), 487-513.

- [15] *H. Federer*, Geometric Measures theory, NewYork, Springer Verlag, 1969.
- [16] *J. Fornæss and N. Sibony*, Oka's inequality for currents and applications, *Math. Ann.*, **301** (1995), 399-419.
- [17] *L. Garnett*, Foliations, the ergodic theorem and Brownian motion, *J. Funct. Anal.*, **51** (1983), no. 3, 285–311.
- [18] *R. Harvey and B. Lawson*, On boundaries of complex analytic varieties I, *Ann. of Math.*, II. Ser., **102** (1975), 223-290.
- [19] *R. Harvey and B. Shiffman*, A characterization of holomorphic chains, *Ann. Math.*, **99** (1974), 553-587.
- [20] *J. King*, The currents defined by analytic varieties, *Acta Math.*, **127** (1971), 185-220.
- [21] *M.G. Lawrence*, Polynomial hulls of rectifiable curves, *Amer. J. Math.*, **117** (1995), 405-417.
- [22] *M.G. Lawrence*, Polynomial hull of sets of finite length in strictly convex domain, *Manuscript* (1997).
- [23] *P. Mattila*, Geometry of Sets and Measures in Euclidean Spaces, *Cambridge University Press*, 1975.
- [24] *B. Shiffman*, Complete characterization of holomorphic chains of codimension one, *Math. Ann.*, **274** (1986), 233-256.
- [25] *N. Sibony*, Quelques problèmes de prolongement de courants en analyse complexe, *Duke Math. J.*, **52** (1985), 157-197.
- [26] *Y.T. Siu*, Analyticity of Sets Associated to Lelong Numbers and the Extension of Closed Positive Currents, *Inventiones math.*, **27** (1974), 53-156.
- [27] *H. Skoda*, Sous-ensemble analytique d'ordre fini ou infini dans $\mathbb{C}^n$, *Bull. Soc. math. France*, **100** (1972), 353-408.
- [28] *H. Skoda*, Prolongement des courants positifs, fermés de masse finie, *Inv. Math.*, **27** (1974), 53-156.

- [29] *G. Stolzenberg*, Uniform approximation on smooth curves, *Acta Math.*, **115** (1966), 185-198.
- [30] *J. Wermer*, The hull of a curve in $\mathbb{C}^n$, *Ann. of Math.*, **68** (1958), 550-561.
- [31] *W.P. Ziemer*, Weakly Differentiable Functions, Springer-Verlag, 1989.

Tien-Cuong Dinh
Mathématique - Bâtiment 425
UMR 8628, Université Paris-Sud
91405 ORSAY Cedex (France)
TienCuong.Dinh@math.u-psud.fr